

Product datasheet for TA350060

HSD17B2 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	WB: 500-2000 WB positive control: Human fetal liver tissue IHC: 50-200 Positive control: Human esophagus cancer Predicted cell location: Cytoplasm
Reactivity:	Human
Host:	Rabbit
lsotype:	lgG
Clonality:	Polyclonal
Immunogen:	Fusion protein of human HSD17B2
Formulation:	pH7.4 PBS, 0.05% NaN3, 40% Glyceroln
Concentration:	lot specific
Purification:	Antigen affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	43 kDa
Gene Name:	hydroxysteroid (17-beta) dehydrogenase 2
Database Link:	<u>NP_002144</u> <u>Entrez Gene 3294 Human</u> <u>P37059</u>

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


	HSD17B2 Rabbit Polyclonal Antibody – TA350060
Background:	17β-HSD2 (17β hydroxysteroid dehydrogenase type 2) belongs to the 17β-HSD family of proteins that regulate the availability of steroids within a tissue. 17β-HSD2 converts active steroids to their inactive form through its oxidative activity. It is a key player in the inactivation of Estradiol and testosterone. Due to the affects that 17β-HSD2 has on the availability of estrogen, it has been extensively investigated for playing a possible role in breast tumor development, colon cancer development and the pathophysiology of endometriosis.
Synonyms:	EDH17B2; HSD17; SDR9C2
Protein Families	: Druggable Genome, Transmembrane
Protein Pathwa	ys: Androgen and estrogen metabolism, Metabolic pathways

Product images:

%// _




Gel: 8%SDS-PAGE Lysate: 40 µg Lane: Human fetal liver tissue Primary antibody: TA350060 (HSD17B2 Antibody) at dilution 1/300 Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution Exposure time: 1 minute



Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using TA350060 (HSD17B2 Antibody) at dilution 1/50 (Original magnification: ×200)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using TA350060 (HSD17B2 Antibody) at dilution 1/50, treated with fusion protein. (Original magnification: ×200)

Immunohistochemistry of paraffin-embedded Human brain tissue using TA350060 (HSD17B2 Antibody) at dilution 1/50 (Original magnification: ×200)

Immunohistochemistry of paraffin-embedded Human brain tissue using TA350060 (HSD17B2 Antibody) at dilution 1/50, treated with fusion protein. (Original magnification: ×200)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US