

Product datasheet for TA350183

MRPL28 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 1000-5000

WB positive control: 293T cells and human testis tissue, A375 cells and human normal kidney

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

tissue

IHC: 100-300

Positive control: Human thyroid cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein of human MRPL28

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 30 kDa

Gene Name: mitochondrial ribosomal protein L28

Database Link: NP 006419

Entrez Gene 68611 MouseEntrez Gene 10573 Human

Q13084






Background:

Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly in sequence, and sometimes in biochemical properties, which prevents easy recognition by sequence homology.

Synonyms: MAAT1; p15

Product images:




Gel: 8%SDS-PAGE Lysate: 40 µg Lane 1-4: 293T cells human testis tissue A375 cells human normal kidney tissue Primary antibody: TA350183 (MRPL28 Antibody) at dilution 1/600

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 2 seconds




Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA350183 (MRPL28 Antibody) at dilution 1/45 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA350183 (MRPL28 Antibody) at dilution 1/45, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using TA350183 (MRPL28 Antibody) at dilution 1/45 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using TA350183 (MRPL28 Antibody) at dilution 1/45, treated with fusion protein. (Original magnification: ×200)