

Product datasheet for TA350222S

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

NDUFV2 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB: 200-1000

WB positive control: Jurkat, Raji and A431 cells, mouse skeletal muscle tissue and NIH/3T3

cells

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein of human NDUFV2

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 27 kDa

Gene Name: NADH:ubiquinone oxidoreductase core subunit V2

Database Link: NP 066552

Entrez Gene 72900 MouseEntrez Gene 81728 RatEntrez Gene 4729 Human

P19404

Background: The NADH-ubiquinone oxidoreductase complex (complex I) of the mitochondrial respiratory

chain catalyzes the transfer of electrons from NADH to ubiquinone, and consists of at least 43 subunits. The complex is located in the inner mitochondrial membrane. This gene encodes the 24 kDa subunit of complex I, and is involved in electron transfer. Mutations in this gene are implicated in Parkinson's disease, bipolar disorder, schizophrenia, and have been found

in one case of early onset hypertrophic cardiomyopathy and encephalopathy. A non-

transcribed pseudogene of this locus is found on chromosome 19.

Synonyms: CI-24k






Protein Families: Druggable Genome

Protein Pathways: Alzheimer's disease, Huntington's disease, Metabolic pathways, Oxidative phosphorylation,

Parkinson's disease

Product images:



Gel: 10%SDS-PAGE Lysate: 40 µg

Lane 1-5: Jurkat cells

Raji cells A431 cells

mouse skeletal muscle tissue

NIH/3T3 cells

Primary antibody: [TA350222] (NDUFV2 Antibody)

at dilution 1/250

Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution

Exposure time: 5 seconds