

Product datasheet for **TA350233S**

NOC2L Rabbit Polyclonal Antibody


Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	WB: 200-1000 WB positive control: Hela cells IHC: 25-100 Positive control: Human prostate cancer Predicted cell location: Cytoplasm or Nucleus
Reactivity:	Human, Mouse
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Fusion protein of human NOC2L
Formulation:	pH7.4 PBS, 0.05% NaN ₃ , 40% Glycerol
Purification:	Antigen affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	85 kDa
Gene Name:	NOC2 like nucleolar associated transcriptional repressor
Database Link:	NP_056473 Entrez Gene 57741 Mouse Entrez Gene 26155 Human Q9Y3T9
Background:	Histone modification by histone acetyltransferases (HAT) and histone deacetylases (HDAC) can control major aspects of transcriptional regulation. NOC2L represents a novel HDAC-independent inhibitor of histone acetyltransferase (INHAT)
Synonyms:	NET7; NET15; NIR; PPP1R112
Protein Families:	Stem cell - Pluripotency




[View online »](#)


Product images:




Gel: 6%SDS-PAGE
Lysate: 40 µg
Lane: HeLa cells
Primary antibody: [TA350233] (NOC2L Antibody)
at dilution 1/200
Secondary antibody: Goat anti rabbit IgG at
1/8000 dilution
Exposure time: 5 seconds




Immunohistochemistry of paraffin-embedded
Human prostate cancer tissue using [TA350233]
(NOC2L Antibody) at dilution 1/20 (Original
magnification: ×200)



Immunohistochemistry of paraffin-embedded Human prostate cancer tissue using [TA350233] (NOC2L Antibody) at dilution 1/20, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human breast cancer tissue using [TA350233] (NOC2L Antibody) at dilution 1/20 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human breast cancer tissue using [TA350233] (NOC2L Antibody) at dilution 1/20, treated with fusion protein. (Original magnification: ×200)