

Product datasheet for TA350490

LEFTY2 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: 293T cells

IHC: 25-100

Positive control: Human thyroid cancer

Predicted cell location: Cytoplasm and Cell membrane

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Full length fusion protein

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 41 kDa

Gene Name: left-right determination factor 2

Database Link: NP 003231

Entrez Gene 320202 MouseEntrez Gene 7044 Human

000292



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:


This gene encodes a member of the TGF-beta family of proteins. The encoded protein is secreted and plays a role in left-right asymmetry determination of organ systems during development. The protein may also play a role in endometrial bleeding. Mutations in this gene have been associated with left-right axis malformations, particularly in the heart and lungs. Some types of infertility have been associated with dysregulated expression of this gene in the endometrium. Alternative processing of this protein can yield three different products. This gene is closely linked to both a related family member and a related pseudogene. Alternate splicing of this gene results in multiple transcript variants.

Synonyms: EBAF; LEFTA; LEFTYA; TGFB4

Protein Families: Druggable Genome, Secreted Protein

Protein Pathways: TGF-beta signaling pathway

Product images:



Gel: 8%SDS-PAGE Lysate: 40 µg Lane: 293T cells


Primary antibody: TA350490 (LEFTY2 Antibody) at

dilution 1/800

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 30 seconds




Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA350490 (LEFTY2 Antibody) at dilution 1/35 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA350490 (LEFTY2 Antibody) at dilution 1/35, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA350490 (LEFTY2 Antibody) at dilution 1/35 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA350490 (LEFTY2 Antibody) at dilution 1/35, treated with fusion protein. (Original magnification: ×200)