

## **Product datasheet for TA350572**

## **UGT1A10** Rabbit Polyclonal Antibody

## **Product data:**

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: WB: 200-1000

WB positive control: A431 cell

Reactivity: Human

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Fusion protein of human UGT1A10

**Formulation:** pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

**Concentration:** lot specific

**Purification:** Antigen affinity purification

**Conjugation:** Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 60 kDa

**Gene Name:** UDP glucuronosyltransferase family 1 member A10

Database Link: NP 061948

Entrez Gene 54575 Human

Q9HAW8

**Background:** This gene encodes a UDP-glucuronosyltransferase, an enzyme of the glucuronidation

pathway that transforms small lipophilic molecules, such as steroids, bilirubin, hormones, and drugs, into water-soluble, excretable metabolites. This gene is part of a complex locus that encodes several UDP-glucuronosyltransferases. The locus includes thirteen unique alternate first exons followed by four common exons. Four of the alternate first exons are

considered pseudogenes.

Synonyms: UDPGT; UGT-1J; UGT1-10; UGT1.10; UGT1J



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




**Protein Families:** 

Transmembrane

**Protein Pathways:** 

Androgen and estrogen metabolism, Ascorbate and aldarate metabolism, Drug metabolism - cytochrome P450, Drug metabolism - other enzymes, Metabolic pathways, Metabolism of xenobiotics by cytochrome P450, Pentose and glucuronate interconversions, Porphyrin and chlorophyll metabolism, Retinol metabolism, Starch and sucrose metabolism

## **Product images:**



Gel: 6%SDS-PAGE Lysate: 40 µg Lane: A431 cell

Primary antibody: TA350572 (UGT1A10 Antibody)

at dilution 1/250

Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution

Exposure time: 10 seconds