

Product datasheet for TA350629S

Androgen Receptor (AR) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 200-1000

WB positive control: LOVO cell lysate

IHC: 50-100

Positive control: Human gastric cancer

Predicted cell location: Nucleus and Cytoplasm

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide of human AR

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: androgen receptor

Database Link: NP 000035

Entrez Gene 367 Human

P10275

Background: The androgen receptor gene is more than 90 kb long and codes for a protein that has 3

major functional domains: the N-terminal domain, DNA-binding domain, and androgen-binding domain. The protein functions as a steroid-hormone activated transcription factor. Upon binding the hormone ligand, the receptor dissociates from accessory proteins, translocates into the nucleus, dimerizes, and then stimulates transcription of androgen responsive genes. This gene contains 2 polymorphic trinucleotide repeat segments that encode polyglutamine and polyglycine tracts in the N-terminal transactivation domain of its

protein.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Synonyms: AIS; AR8; DHTR; HUMARA; HYSP1; KD; NR3C4; SBMA; SMAX1; TFM

Protein Families: Druggable Genome, Nuclear Hormone Receptor, Transcription Factors

Protein Pathways: Oocyte meiosis, Pathways in cancer, Prostate cancer

Product images:



Gel: 6%SDS-PAGE Lysate: 40 µg

Lane: LOVO cell lysate


Primary antibody: [TA350629] (AR Antibody) at

dilution 1/200

Secondary antibody: Goat anti rabbit IgG at


1/5000 dilution

Exposure time: 40 seconds



Immunohistochemistry of paraffin-embedded Human gastric cancer tissue using [TA350629] (AR Antibody) at dilution 1/35 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human gastric cancer tissue using [TA350629] (AR Antibody) at dilution 1/35, treated with synthetic peptide. (Original magnification: ×200)