

Product datasheet for TA350640

Insulin (INS) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC

Recommended Dilution: IHC: 25-100

Positive control: Human esophagus cancer

Predicted cell location: Cytoplasm

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide of human INS

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: insulin

Database Link: NP 000198

Entrez Gene 3630 Human

P01308

Background: After removal of the precursor signal peptide, proinsulin is post-translationally cleaved into

three peptides: the B chain and A chain peptides, which are covalently linked via two disulfide

bonds to form insulin, and C-peptide. Binding of insulin to the insulin receptor (INSR)

stimulates glucose uptake. A multitude of mutant alleles with phenotypic effects have been identified. There is a read-through gene, INS-IGF2, which overlaps with this gene at the 5' region and with the IGF2 gene at the 3' region. Alternative splicing results in multiple

transcript variants.

Synonyms: IDDM; IDDM1; IDDM2; ILPR; IRDN; MODY10



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Protein Families: Druggable Genome, ES Cell Differentiation/IPS, Secreted Protein


Protein Pathways: Insulin signaling pathway, Maturity onset diabetes of the young, mTOR signaling pathway,
Oocyte meiosis, Progesterone-mediated oocyte maturation, Prostate cancer, Regulation of

actin cytoskeleton, Regulation of autophagy, Type I diabetes mellitus, Type II diabetes mellitus

Product images:




Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using TA350640 (INS Antibody) at dilution 1/35 (Original magnification: ×200)




Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using TA350640 (INS Antibody) at dilution 1/35, treated with synthetic peptide. (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA350640 (INS Antibody) at dilution 1/35 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA350640 (INS Antibody) at dilution 1/35, treated with synthetic peptide. (Original magnification: ×200)