

# **Product datasheet for TA350655**

## EAAT3 (SLC1A1) Rabbit Polyclonal Antibody

### **Product data:**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	WB: 500-2000 WB positive control: Human placenta tissue and normal liver tissue IHC: 25-100 Positive control: Human liver cancer Predicted cell location: Cell membrane and Cytoplasm
Reactivity:	Human
Host:	Rabbit
lsotype:	IgG
Clonality:	Polyclonal
Immunogen:	Synthetic peptide of human SLC1A1
Formulation:	pH7.4 PBS, 0.05% NaN3, 40% Glyceroln
Concentration:	lot specific
Purification:	Antigen affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	57 kDa
Gene Name:	solute carrier family 1 member 1
Database Link:	<u>NP_004161</u> <u>Entrez Gene 6505 Human</u> <u>P43005</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	EAAT3 (SLC1A1) Rabbit Polyclonal Antibody – TA350655
Background:	This gene encodes a member of the high-affinity glutamate transporters that play an essential role in transporting glutamate across plasma membranes. In brain, these transporters are crucial in terminating the postsynaptic action of the neurotransmitter glutamate, and in maintaining extracellular glutamate concentrations below neurotoxic levels. This transporter also transports aspartate, and mutations in this gene are thought to cause dicarboxylicamino aciduria, also known as glutamate-aspartate transport defect.
Synonyms:	DCBXA; EAAC1; EAAT3; SCZD18
Protein Families	Druggable Genome, Transmembrane

# **Product images:**




Gel: 8%SDS-PAGE Lysate: 40 μg Lane 1-2: Human placenta tissue and normal liver tissue Primary antibody: TA350655 (SLC1A1 Antibody) at dilution 1/400 Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution Exposure time: 4 minutes



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA350655 (SLC1A1 Antibody) at dilution 1/30 (Original magnification: ×200)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA350655 (SLC1A1 Antibody) at dilution 1/30, treated with synthetic peptide. (Original magnification: ×200)

Immunohistochemistry of paraffin-embedded Human breast cancer tissue using TA350655 (SLC1A1 Antibody) at dilution 1/30 (Original magnification: ×200)

Immunohistochemistry of paraffin-embedded Human breast cancer tissue using TA350655 (SLC1A1 Antibody) at dilution 1/30, treated with synthetic peptide. (Original magnification: ×200)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US