

Product datasheet for TA350833

PAK3 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 200-1000

WB positive control: Mouse brain tissue

IHC: 20-100

Positive control: Human esophagus cancer

Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide of human PAK3

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 62 kDa

Gene Name: p21 (RAC1) activated kinase 3

Database Link: NP 002569

Entrez Gene 18481 MouseEntrez Gene 29433 RatEntrez Gene 5063 Human

<u>075914</u>



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background: PAK proteins are critical effectors that link Rho GTPases to cytoskeleton reorganization and

nuclear signaling. PAK proteins, a family of serine/threonine p21-activating kinases, serve as targets for the small GTP binding proteins Cdc42 and RAC and have been implicated in a wide range of biological activities. The protein encoded by this gene forms an activated complex with GTP-bound RAS-like (P21), CDC2 and RAC1 proteins which then catalyzes a variety of

targets.


Synonyms: ARA; beta-PAK; bPAK; MRX30; MRX47; OPHN3; PAK-3; PAK3beta

Protein Families: Druggable Genome, Protein Kinase, Stem cell - Pluripotency

Protein Pathways: Axon guidance, ErbB signaling pathway, Focal adhesion, Regulation of actin cytoskeleton,

Renal cell carcinoma, T cell receptor signaling pathway

Product images:



Gel: 6%SDS-PAGE Lysate: 40 μg

Lane: Mouse brain tissue


Primary antibody: TA350833 (PAK3 Antibody) at

dilution 1/200

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 1 minute




Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using TA350833 (PAK3 Antibody) at dilution 1/25 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using TA350833 (PAK3 Antibody) at dilution 1/25, treated with synthetic peptide. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA350833 (PAK3 Antibody) at dilution 1/25 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA350833 (PAK3 Antibody) at dilution 1/25, treated with synthetic peptide. (Original magnification: ×200)