

Product datasheet for TA350872S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ARIH2 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB: 500-2000

WB positive control: Hela, Jurkat, 293T and K562 cells, mouse testis tissue

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide of human ARIH2

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 58 kDa

Gene Name: ariadne RBR E3 ubiquitin protein ligase 2

Database Link: NP 006312

Entrez Gene 23807 MouseEntrez Gene 10425 Human

O95376

Background: TRIAD1, also known as ARIH2 (ariadne homolog 2) or ARI2, is a 493 amino acid protein that

contains one IBR-type zinc finger and two RING-type zinc fingers and belongs to the ariadne subfamily of RBR proteins. Localized to the nucleus, TRIAD1 interacts with UBE2L3 and is thought to act as an E3 ubiquitin-protein ligase, functioning to accept ubiquitin from E2 ubiquitin-conjugating enzymes and transfer the acquired ubiquitin residue to target

substrates. TRIAD1 is subject to post-translational DNA damage-dependent phosphorylation, probably by ATM or ATR. The gene encoding TRIAD1 maps to human chromosome 3, which houses over 1,100 genes, including a chemokine receptor (CKR) gene cluster and a variety of

human cancer-related gene loci.






Synonyms: ARI2; TRIAD1

Protein Families: Druggable Genome

Product images:



Gel: 8%SDS-PAGE Lysate: 40 µg Lane 1-5: Hela cells Jurkat cells 293T cells K562 cells

mouse testis tissue

Primary antibody: [TA350872] (ARIH2 Antibody) at

dilution 1/550

Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution

Exposure time: 40 seconds