

Product datasheet for TA350892S

Rex1 (ZFP42) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB: 500-2000

WB positive control: A549, K562 and PC3 cells

Reactivity: Human

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide of human ZFP42

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 35 kDa

Gene Name: ZFP42 zinc finger protein

Database Link: NP 777560

Entrez Gene 132625 Human

Q96MM3

Background: Rex-1 (for reduced expression), also designated zinc finger protein 42 (ZFP42), is an acidic zinc

finger protein. Rex-1 contains four repeats of the zinc finger nucleic acid-binding motif and a potential acidic activator domain, suggesting that it is a regulatory protein. Rex-1 localizes to the nucleus and is highly expressed in embryonic stem (ES) and undifferentiated murine F9 teratocarcinoma cells. At the transcriptional level, expression of Rex-1 is reduced when F9 cells are induced to differentiate by the addition of retinoic acid (RA), and Rex-1 repression is enhanced by E1A. The Oct-3/4 transcription factor can either activate or repress the Rex-1 promoter, depending on the cellular environment, while Oct-6 can lower the expression of

Rex-1.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms: REX-1; REX1; zfp-42; ZNF754

Protein Families: Adult stem cells, Embryonic stem cells, ES Cell Differentiation/IPS, Stem cell - Pluripotency

Product images:



Gel: 12%SDS-PAGE Lysate: 40 µg Lane 1-3: A549 cells

K562 cells PC3 cells

Primary antibody: [TA350892] (ZFP42 Antibody) at

dilution 1/500

Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution

Exposure time: 30 seconds