

Recommended Dilution: WB: 500-2000

# Product datasheet for TA350933S

# NDNL2 (NSMCE3) Rabbit Polyclonal Antibody

IHC, WB

IHC: 50-200

Human

Rabbit

Polyclonal

Unconjugated

34 kDa

NP 619649

Q96MG7

lgG

**Primary Antibodies** 

WB positive control: LO2 cell lysate

Synthetic peptide of human NDNL2

Antigen affinity purification

Store at -20°C as received.

Entrez Gene 56160 Human

pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

Stable for 12 months from date of receipt.

NSE3 homolog, SMC5-SMC6 complex component

Positive control: Human colorectal cancer Predicted cell location: Cytoplasm and Nucleus

## **Product data:**

**Product Type:** 

**Applications:** 

**Reactivity:** 

Host:

Isotype:

**Clonality:** 

Immunogen:

Formulation:

**Purification:** 

**Conjugation:** 

Gene Name:

Database Link:

**Predicted Protein Size:** 

Storage: Stability:

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


### **ORIGENE** NDNL2 (NSMCE3) Rabbit Polyclonal Antibody – TA350933S

#### Background:


The protein encoded by this gene is part of the SMC5-6 chromatin reorganizing complex and is a member of the MAGE superfamily. This is an intronless gene. Component of the SMC5-SMC6 complex, a complex involved in repair of DNA double-strand breaks by homologous recombination. The complex may promote sister chromatid homologous recombination by recruiting the SMC1-SMC3 cohesin complex to double-strand breaks. The complex is required for telomere maintenance via recombination in ALT (alternative lengthening of telomeres) cell lines and mediates sumoylation of shelterin complex (telosome) components which is proposed to lead to shelterin complex disassembly in ALT-associated PML bodies (APBs).

Synonyms: HCA4; MAGEG1; MAGEL3; NSE3; NSMCE3

### **Product images:**




Gel: 8%SDS-PAGE Lysate: 40 µg Lane: LO2 cell lysate Primary antibody: [TA350933] (NSMCE3 Antibody) at dilution 1/400 Secondary antibody: Goat anti rabbit lgG at 1/5000 dilution Exposure time: 90 seconds



Immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using [TA350933] (NSMCE3 Antibody) at dilution 1/55 (Original magnification: ×200)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using [TA350933] (NSMCE3 Antibody) at dilution 1/55, treated with synthetic peptide. (Original magnification: ×200)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US