

### **Product datasheet for TA350952S**

# Caveolin 3 (CAV3) Rabbit Polyclonal Antibody

#### **Product data:**

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Human fetal muscle tissue

IHC: 50-200

Positive control: Human gasrtic cancer Predicted cell location: Cytoplasm

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Synthetic peptide of human CAV3

**Formulation:** pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 17 kDa

Gene Name: caveolin 3

Database Link: NP 203123

Entrez Gene 859 Human

P56539

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn





Background:


This gene encodes a caveolin family member, which functions as a component of the caveolae plasma membranes found in most cell types. Caveolin proteins are proposed to be scaffolding proteins for organizing and concentrating certain caveolin-interacting molecules. Mutations identified in this gene lead to interference with protein oligomerization or intracellular routing, disrupting caveolae formation and resulting in Limb-Girdle muscular dystrophy type-1C (LGMD-1C), hyperCKemia or rippling muscle disease (RMD). Alternative splicing has been identified for this locus, with inclusion or exclusion of a differentially spliced intron. In addition, transcripts utilize multiple polyA sites and contain two potential translation initiation sites.

Synonyms: LGMD1C; LQT9; VIP-21; VIP21

**Protein Families:** Druggable Genome, Transmembrane

**Protein Pathways:** Focal adhesion

# **Product images:**



Gel: 10%SDS-PAGE Lysate: 40 μg

Lane: Human fetal muscle tissue

Primary antibody: [TA350952] (CAV3 Antibody) at


dilution 1/550

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 15 seconds






Immunohistochemistry of paraffin-embedded Human gasrtic cancer tissue using [TA350952] (CAV3 Antibody) at dilution 1/40 (Original magnification: ×200)




Immunohistochemistry of paraffin-embedded Human gasrtic cancer tissue using [TA350952] (CAV3 Antibody) at dilution 1/40, treated with synthetic peptide. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA350952] (CAV3 Antibody) at dilution 1/40 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA350952] (CAV3 Antibody) at dilution 1/40, treated with synthetic peptide. (Original magnification: ×200)