

## **Product datasheet for TA350988S**

## **APOL6 Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

Applications: IHC

Recommended Dilution: IHC: 50-200

Positive control: Human tonsil Predicted cell location: Cytoplasm

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide of human APOL6

**Formulation:** pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

**Purification:** Antigen affinity purification

**Conjugation:** Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Gene Name:** apolipoprotein L6

Database Link: NP 085144

Entrez Gene 80830 Human

Q9BWW8

**Background:** This gene is a member of the apolipoprotein L gene family. The encoded protein is found in

the cytoplasm, where it may affect the movement of lipids or allow the binding of lipids to organelles. Widely expressed; highly expressed in the uterus, fetal brain and spinal cord, also detected in heart, liver, lung, colon, spleen, thymus, prostate, placenta, adrenal gland, salivary

and mammary gland.

Synonyms: APOL-VI; APOLVI




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Product images:**




Immunohistochemistry of paraffin-embedded Human tonsil tissue using [TA350988] (APOL6 Antibody) at dilution 1/60 (Original magnification: ×200)




Immunohistochemistry of paraffin-embedded Human tonsil tissue using [TA350988] (APOL6 Antibody) at dilution 1/60, treated with synthetic peptide. (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA350988] (APOL6 Antibody) at dilution 1/60 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA350988] (APOL6 Antibody) at dilution 1/60, treated with synthetic peptide. (Original magnification: ×200)