

Product datasheet for TA351007S

BLNK Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Mouse liver, kidney and pancreas tissue

IHC: 50-200

Positive control: Human lung cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide of human BLNK

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 50 kDa

Gene Name: B-cell linker

Database Link: NP 037446

Entrez Gene 17060 MouseEntrez Gene 499356 RatEntrez Gene 29760 Human

Q8WV28



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:


This gene encodes a cytoplasmic linker or adaptor protein that plays a critical role in B cell development. This protein bridges B cell receptor-associated kinase activation with downstream signaling pathways, thereby affecting various biological functions. The phosphorylation of five tyrosine residues is necessary for this protein to nucleate distinct signaling effectors following B cell receptor activation. Mutations in this gene cause hypoglobulinemia and absent B cells, a disease in which the pro- to pre-B-cell transition is developmentally blocked. Deficiency in this protein has also been shown in some cases of pre-B acute lymphoblastic leukemia.

Synonyms: AGM4; BASH; bca; BLNK-S; LY57; SLP-65; SLP65

Protein Families: Druggable Genome

Protein Pathways: B cell receptor signaling pathway, Primary immunodeficiency

Product images:



Gel: 8%SDS-PAGE Lysate: 40 µg Lane 1-3: Mouse liver Mouse kidney tissue Mouse pancreas tissue


Primary antibody: [TA351007] (BLNK Antibody) at

dilution 1/550

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 40 seconds




Immunohistochemistry of paraffin-embedded Human lung cancer tissue using [TA351007] (BLNK Antibody) at dilution 1/50 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human lung cancer tissue using [TA351007] (BLNK Antibody) at dilution 1/50, treated with synthetic peptide. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human colon cancer tissue using [TA351007] (BLNK Antibody) at dilution 1/50 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human colon cancer tissue using [TA351007] (BLNK Antibody) at dilution 1/50, treated with synthetic peptide. (Original magnification: ×200)