

Product datasheet for TA351017

CBL Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB: 500-2000

WB positive control: 231 cells

Reactivity: Human, Mouse

Host: Rabbit Isotype: **IgG**

Clonality:

Polyclonal

Synthetic peptide of human CBL Immunogen:

pH7.4 PBS, 0.05% NaN3, 40% Glyceroln Formulation:

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated

Store at -20°C as received. Storage:

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 100 kDa

Gene Name: Cbl proto-oncogene

Database Link: NP 005179

Entrez Gene 12402 MouseEntrez Gene 867 Human

P22681



9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn





Background:

This gene is a proto-oncogene that encodes a RING finger E3 ubiquitin ligase. The encoded protein is one of the enzymes required for targeting substrates for degradation by the proteasome. This protein mediates the transfer of ubiquitin from ubiquitin conjugating enzymes (E2) to specific substrates. This protein also contains an N-terminal phosphotyrosine binding domain that allows it to interact with numerous tyrosine-phosphorylated substrates and target them for proteasome degradation. As such it functions as a negative regulator of many signal transduction pathways. This gene has been found to be mutated or translocated in many cancers including acute myeloid leukaemia. Mutations in this gene are also the cause of Noonan syndrome-like disorder

Synonyms: C-CBL; CBL2; FRA11B; NSLL; RNF55

Protein Families: Druggable Genome, Transcription Factors

Protein Pathways: Chronic myeloid leukemia, Endocytosis, ErbB signaling pathway, Insulin signaling pathway,

Jak-STAT signaling pathway, Pathways in cancer, T cell receptor signaling pathway, Ubiquitin

mediated proteolysis

Product images:



Gel: 10%SDS-PAGE Lysate: 40 µg Lane: 231 cells

Primary antibody: TA351017 (CBL Antibody) at

dilution 1/750

Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution

Exposure time: 1 minute