

## **Product datasheet for TA351025S**

## 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

OriGene Technologies, Inc.

EU: info-de@origene.com CN: techsupport@origene.cn

## Cell adhesion molecule 4 (CADM4) Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 200-1000

WB positive control: RAW264.7 cells and mouse brain tissue

IHC: 25-100

Positive control: Human liver cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Synthetic peptide of human CADM4

**Formulation:** pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 43 kDa

**Gene Name:** cell adhesion molecule 4

Database Link: NP 660339

Entrez Gene 260299 MouseEntrez Gene 365216 RatEntrez Gene 199731 Human

Q8NFZ8






Background:

IGSF4C (immunoglobulin superfamily member 4C), also known as CADM4 (cell adhesion molecule 4), NECL4 or TSLL2, is a 388 amino acid single-pass type I membrane protein that contains one Ig-like V-type domain and two Ig-like C2-type domains. Expressed in kidney, brain and prostate, IGSF4C exists as both a monomer and a homodimer and is thought to be involved in cell-cell adhesion, possibly playing a role in tumor suppression. Human IGSF4C shares 98% sequence identity with its mouse counterpart, suggesting a conserved role between species. The gene encoding IGSF4C maps to human chromosome 19, which is the genetic home for a number of immunoglobulin superfamily members, including the killer cell and leukocyte Ig-like receptors, a number of ICAMs, the CEACAM and PSG family and Fc receptors (FcRs)

Synonyms: IGSF4C; Necl-4; NECL4; synCAM4; TSLL2

**Protein Families:** Transmembrane

## **Product images:**




Gel: 10%SDS-PAGE Lysate: 40 µg Lane 1-2: RAW264.7 cells

mouse brain tissue Primary antibody: [TA351025] (CADM4 Antibody) at dilution 1/300

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 20 seconds




Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA351025] (CADM4 Antibody) at dilution 1/25 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA351025] (CADM4 Antibody) at dilution 1/25, treated with synthetic peptide. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human ovarian cancer tissue using [TA351025] (CADM4 Antibody) at dilution 1/25 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human ovarian cancer tissue using [TA351025] (CADM4 Antibody) at dilution 1/25, treated with synthetic peptide. (Original magnification: ×200)