

# Product datasheet for TA351048

## **CD36 Rabbit Polyclonal Antibody**

### **Product data:**

#### **Product Type: Primary Antibodies Applications:** IHC, WB Recommended Dilution: WB: 500-2000 WB positive control: Human placenta and fat tissue IHC: 50-200 Positive control: Human thyroid cancer Predicted cell location: Cytoplasm **Reactivity:** Human Host: Rabbit Isotype: lgG **Clonality:** Polyclonal Immunogen: Synthetic peptide of human CD36 pH7.4 PBS, 0.05% NaN3, 40% Glyceroln Formulation: Concentration: lot specific **Purification:** Antigen affinity purification **Conjugation:** Unconjugated Storage: Store at -20°C as received. Stability: Stable for 12 months from date of receipt. Predicted Protein Size: 53 kDa Gene Name: CD36 molecule Database Link: NP 000063 Entrez Gene 948 Human P16671

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


### **CD36** Rabbit Polyclonal Antibody – TA351048

Background:	The protein encoded by this gene is the fourth major glycoprotein of the platelet surface and serves as a receptor for thrombospondin in platelets and various cell lines. Since thrombospondins are widely distributed proteins involved in a variety of adhesive processes, this protein may have important functions as a cell adhesion molecule. It binds to collagen, thrombospondin, anionic phospholipids and oxidized LDL. It directly mediates cytoadherence of Plasmodium falciparum parasitized erythrocytes and it binds long chain fatty acids and may function in the transport and/or as a regulator of fatty acid transport.
Synonyms:	BDPLT10; CHDS7; FAT; GP3B; GP4; GPIV; PASIV; SCARB3
Protein Families:	Druggable Genome, Transmembrane
Protein Pathways:	Adipocytokine signaling pathway, ECM-receptor interaction, Hematopoietic cell lineage, PPAR signaling pathway

### **Product images:**




Gel: 8%SDS-PAGE Lysate: 40 µg Lane 1-2: Human placenta tissue Human fat tissue Primary antibody: TA351048 (CD36 Antibody) at dilution 1/1000 Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution Exposure time: 1 minute



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA351048 (CD36 Antibody) at dilution 1/60 (Original magnification: ×200)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA351048 (CD36 Antibody) at dilution 1/60, treated with synthetic peptide. (Original magnification: ×200)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US