

Product datasheet for **TA351077S**

CHRNA2 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	WB: 500-2000 WB positive control: A172 and A549 cells IHC: 25-100 Positive control: Human cervical cancer Predicted cell location: Cytoplasm
Reactivity:	Human
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Synthetic peptide of human CHRNA2
Formulation:	pH7.4 PBS, 0.05% NaN ₃ , 40% Glycerol
Purification:	Antigen affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	60 kDa
Gene Name:	cholinergic receptor nicotinic alpha 2 subunit
Database Link:	NP_000733 Entrez Gene 1135 Human Q15822



[View online »](#)

Background:

Nicotinic acetylcholine receptors (nAChRs) are ligand-gated ion channels formed by a pentameric arrangement of alpha and beta subunits to create distinct muscle and neuronal receptors. Neuronal receptors are found throughout the peripheral and central nervous system where they are involved in fast synaptic transmission. This gene encodes an alpha subunit that is widely expressed in the brain. The proposed structure for nAChR subunits is a conserved N-terminal extracellular domain followed by three conserved transmembrane domains, a variable cytoplasmic loop, a fourth conserved transmembrane domain, and a short C-terminal extracellular region. Mutations in this gene cause autosomal dominant nocturnal frontal lobe epilepsy type 4. Single nucleotide polymorphisms (SNPs) in this gene have been associated with nicotine dependence.

Synonyms:


alpha 2 (neuronal); alpha polypeptide 2 (neuronal); cholinergic receptor; nicotinic; OTTHUMP00000128276

Protein Families:


Druggable Genome, Ion Channels: Cys-loop Receptors, Transmembrane

Product images:


Gel: 8%SDS-PAGE
Lysate: 40 µg
Lane 1-2: A172 cells
A549 cells
Primary antibody: [TA351077] (CHRNA2 Antibody)
at dilution 1/950
Secondary antibody: Goat anti rabbit IgG at
1/8000 dilution
Exposure time: 1 minute




Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using [TA351077] (CHRNA2 Antibody) at dilution 1/40 (Original magnification: $\times 200$)



Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using [TA351077] (CHRNA2 Antibody) at dilution 1/40, treated with synthetic peptide. (Original magnification: $\times 200$)



Immunohistochemistry of paraffin-embedded Human brain tissue using [TA351077] (CHRNA2 Antibody) at dilution 1/40 (Original magnification: $\times 200$)



Immunohistochemistry of paraffin-embedded Human brain tissue using [TA351077] (CHRNA2 Antibody) at dilution 1/40, treated with synthetic peptide. (Original magnification: $\times 200$)