

# **Product datasheet for TA351088**

# CMG1 (IFT74) Rabbit Polyclonal Antibody

### **Product data:**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB: 500-2000 WB positive control: HepG2 and 293T cells
Reactivity:	Human
Host:	Rabbit
lsotype:	lgG
Clonality:	Polyclonal
Immunogen:	Synthetic peptide of human IFT74
Formulation:	pH7.4 PBS, 0.05% NaN3, 40% Glyceroln
Concentration:	lot specific
Purification:	Antigen affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	69 kDa
Gene Name:	intraflagellar transport 74
Database Link:	<u>NP_079379</u> <u>Entrez Gene 80173 Human</u> <u>Q96LB3</u>
Background:	Capillary morphogenesis protein 1 (CMG1), also known as intraflagellar transport protein 74 (ITP74) or coiled-coil domain-containing protein 2 (CCDC2), is a 600 amino acid human homologue of IFT-71, a complex B protein supporting intraflagellar transport (IFT) in Chlamydomonas. CMG1 localizes to the cytoplasmic vesicle and is highly expressed in adult and fetal kidney and testis, with lower levels of expression in adult heart, placenta, lung, liver and pancreas, and in fetal heart, lung and liver. CMG1 has been suggested to have a role in the primary cilia of HUVEC, and it also functions as a transcriptional regulator of cyclin D2 in




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

spermatocyte-derived cells.

Synonyms: BBS20; CCDC2; CMG-1; CMG1

# **Product images:**



Gel: 8%SDS-PAGE Lysate: 40 µg Lane 1-2: HepG2 cells 293T cells Primary antibody: TA351088 (IFT74 Antibody) at dilution 1/1400 Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution Exposure time: 15seconds

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US