

# **Product datasheet for TA351198**

## **FPR3 Rabbit Polyclonal Antibody**

### **Product data:**

**Product Type:** Primary Antibodies

Applications: IHC

Recommended Dilution: IHC: 25-100

Positive control: Human colon cancer Predicted cell location: Cytoplasm

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Synthetic peptide of human FPR3

**Formulation:** pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

**Concentration:** lot specific

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Gene Name:** formyl peptide receptor 3

Database Link: NP 002021

Entrez Gene 2359 Human

P25089

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn





Background:


The N-formyl peptide receptor (FPR) family is comprised of three members, FPR, FPR3 (also designated FPRL1, lipoxin A4 receptor, FPRH1 or FPR2) and FPR like-2 (FPRL2), all of which are chemotactic G protein-coupled receptors that contain seven transmembrane domains. These receptors are found on the surface of phagocytic leukocytes, such as neutrophils and monocytes, and each family member contains specific residues, which are responsible for determining its ligand specificity. FPR3 is a promiscuous receptor that binds to several ligands, including lipoxin A4, N-formyl-methionyl-leucyl-phenylalanine (fMLP), serum amyloid A (SAA), prion peptide and the 42 amino acid form of beta amyloid. Upon activation, FPR3 induces migration and calcium mobilization in human monocytes and neutrophils and is involved in inflammatory and host defense responses. FPR3 may mediate inflammation in prion and Alzheimer?s diseases, which makes it a potential target for therapeutic agents.

Synonyms: FML2\_HUMAN; FMLP-R-II; FMLPY; FPRH1; FPRH2; FPRL2; RMLP-R-I

**Protein Families:** Druggable Genome, GPCR, Transmembrane


**Protein Pathways:** Neuroactive ligand-receptor interaction

## **Product images:**




Immunohistochemistry of paraffin-embedded Human colon cancer tissue using TA351198 (FPR3 Antibody) at dilution 1/20 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human colon cancer tissue using TA351198 (FPR3 Antibody) at dilution 1/20, treated with synthetic peptide. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human tonsil tissue using TA351198 (FPR3 Antibody) at dilution 1/20 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human tonsil tissue using TA351198 (FPR3 Antibody) at dilution 1/20, treated with synthetic peptide. (Original magnification: ×200)