

## **Product datasheet for TA351248S**

# **H2AC4** Rabbit Polyclonal Antibody

#### **Product data:**

**Product Type:** Primary Antibodies

**Applications:** IHC, WB

Recommended Dilution: WB: 200-1000

WB positive control: Hela, 231 and K562 cells, human fetal brain tissue

IHC: 50-200

Positive control: Human cervical cancer

Predicted cell location: Nucleus

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen:Synthetic peptide of human HIST1H2ABFormulation:pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 14 kDa

**Gene Name:** histone cluster 1, H2ab

Database Link: NP 003504

Entrez Gene 8335 Human

P04908



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




#### Background:


Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped around a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a member of the histone H2A family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6p22-p21.3.


Synonyms: H2A; H2AFM; m

**Protein Pathways:** Systemic lupus erythematosus

### **Product images:**







human fetal brain tissue


Primary antibody: [TA351248] (H2AC4 Antibody)

at dilution 1/250

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 5 minutes




Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using [TA351248] (H2AC4 Antibody) at dilution 1/30 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using [TA351248] (H2AC4 Antibody) at dilution 1/30, treated with synthetic peptide. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human colon cancer tissue using [TA351248] (H2AC4 Antibody) at dilution 1/30 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human colon cancer tissue using [TA351248] (H2AC4 Antibody) at dilution 1/30, treated with synthetic peptide. (Original magnification: ×200)