

Product datasheet for TA351254S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

HNRPH1 (HNRNPH1) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB: 500-2000

WB positive control: LoVo and A375 cells, mouse brain tissue, HepG2, A431 and Hela cells

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen:Synthetic peptide of human HNRNPH1Formulation:pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 49 kDa

Gene Name: heterogeneous nuclear ribonucleoprotein H1 (H)

Database Link: NP 005511

Entrez Gene 59013 MouseEntrez Gene 140931 RatEntrez Gene 3187 Human

P31943

Background: This gene encodes a member of a subfamily of ubiquitously expressed heterogeneous

nuclear ribonucleoproteins (hnRNPs). The hnRNPs are RNA binding proteins that complex with heterogeneous nuclear RNA. These proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA

metabolism and transport. While all of the hnRNPs are present in the nucleus, some may shuttle between the nucleus and the cytoplasm. The hnRNP proteins have distinct nucleic acid binding properties. The protein encoded by this gene has three repeats of quasi-RRM domains that bind to RNA and is very similar to the family member HNRPF. This gene may be


associated with hereditary lymphedema type I.





Synonyms: hnRNPH; HNRPH; HNRPH1

Product images:



Gel: 8%SDS-PAGE
Lysate: 40 µg
Lane 1-6: LoVo cells
A375 cells
mouse brain tissue
HepG2 cells
A431 cells
Hela cells
Primary antibody: [TA351254] (HNRNPH1
Antibody) at dilution 1/350
Secondary antibody: Goat anti rabbit IgG at
1/8000 dilution

Exposure time: 40 seconds