

Product datasheet for TA351268S

HSD17B1 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Human placenta tissue

IHC: 50-200

Positive control: Human thyroid cancer Predicted cell location: Cytoplasm

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

Immunogen:Synthetic peptide of human HSD17B1Formulation:pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 35 kDa

Gene Name: hydroxysteroid (17-beta) dehydrogenase 1

Database Link: NP 000404

Entrez Gene 3292 Human

P14061

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn





Background: Hydroxysteroid (17-beta) dehydrogenase 1 catalyzes the final step in the formation of

estradiol and testosterone from estrone and androstenedione, respectively. Ovarian granulosa cells and breast tissue both express HSD17B1. Other tissues that express HSD17B1 include testis, placenta, uterus, prostate and adipose tissue. HSD17B1 functions as a

homodimer and prefers NADP(H) over NAD(H) for oxidation and reduction. The gene encoding human HSD17B1 maps to chromosome 17q12-q21. The importance of HSD17B1 to estradiol production suggests the specific inhibition of HSD17B1 may aid in breast cancer


therapy.

Synonyms: EDH17B2; EDHB17; HSD17; SDR28C1

Protein Families: Druggable Genome

Protein Pathways: Androgen and estrogen metabolism, Metabolic pathways

Product images:



Gel: 8%SDS-PAGE Lysate: 40 μg

Lane: Human placenta tissue


Primary antibody: [TA351268] (HSD17B1

Antibody) at dilution 1/700

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 1 second




Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA351268] (HSD17B1 Antibody) at dilution 1/40 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA351268] (HSD17B1 Antibody) at dilution 1/40, treated with synthetic peptide. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human brain tissue using [TA351268] (HSD17B1 Antibody) at dilution 1/40 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human brain tissue using [TA351268] (HSD17B1 Antibody) at dilution 1/40, treated with synthetic peptide. (Original magnification: ×200)