

## **Product datasheet for TA351316**

## **KCNG3 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** IHC, WB

Recommended Dilution: WB: 200-1000

WB positive control: Human breast infiltrative duct tissue

IHC: 10-50

Positive control: Human liver cancer Predicted cell location: Cell membrane

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide of human KCNG3

**Formulation:** pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

**Concentration:** lot specific

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 50 kDa

**Gene Name:** potassium voltage-gated channel modifier subfamily G member 3

Database Link: NP 758847

Entrez Gene 171011 RatEntrez Gene 225030 MouseEntrez Gene 170850 Human

Q8TAE7



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

Voltage-gated potassium (Kv) channels represent the most complex class of voltage-gated ion channels from both functional and structural standpoints. Their diverse functions include regulating neurotransmitter release, heart rate, insulin secretion, neuronal excitability, epithelial electrolyte transport, smooth muscle contraction, and cell volume. This gene encodes a member of the potassium channel, voltage-gated, subfamily G. This member is a gamma subunit functioning as a modulatory molecule.

Synonyms: KV6.3; KV10.1

**Protein Families:** Druggable Genome, Ion Channels: Other, Transmembrane

## **Product images:**



Gel: 8%SDS-PAGE Lysate: 40 μg


Lane: Human breast infiltrative duct tissue Primary antibody: TA351316 (KCNG3 Antibody) at

dilution 1/200

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 30 seconds




Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA351316 (KCNG3 Antibody) at dilution 1/25 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA351316 (KCNG3 Antibody) at dilution 1/25, treated with synthetic peptide. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human breast cancer tissue using TA351316 (KCNG3 Antibody) at dilution 1/25 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human breast cancer tissue using TA351316 (KCNG3 Antibody) at dilution 1/25, treated with synthetic peptide. (Original magnification: ×200)