

Product datasheet for TA351326S

KIDINS220 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB: 200-1000

WB positive control: Raji and Hela cells

Reactivity: Human, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide of human KIDINS220 Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 197 kDa

Gene Name: kinase D-interacting substrate 220kDa

Database Link: NP 065789

Entrez Gene 116478 RatEntrez Gene 57498 Human

Q9ULH0

Background: Ankyrin repeat-rich membrane-spanning protein (ARMS), also designated kinase D-interacting

substance 220 or Kidins220, is a highly conserved protein containing multiple domains, including four putative transmembrane domains and several ankyrin repeats. ARMS is expressed in regions rich in neurotrophin (Trk) and ephrin (Eph) receptors, such as the brain and neuroendocrine cells (where it concentrates at the tip of neurites) and in plastic areas of the adult brain. It is also detected in peripheral blood immature dendritic cells and PC12 cells. ARMS functions as a substrate for protein kinase D and is a downstream target for both Trk and Eph receptors. It is a highly conserved protein, which suggests it has an evolutionary

conserved role. The gene encoding for the protein maps to chromosome 2p24.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Synonyms: ARMS

Protein Families: Druggable Genome, Transmembrane

Protein Pathways: Neurotrophin signaling pathway

Product images:



Gel: 6%SDS-PAGE Lysate: 40 μg Lane 1-2: Raji cells

Hela cells

Primary antibody: [TA351326] (KIDINS220

Antibody) at dilution 1/200

Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution

Exposure time: 20 seconds