

## **Product datasheet for TA351364**

## **LPP Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

**Applications:** IHC, WB

Recommended Dilution: WB: 200-1000

WB positive control: HepG2 cel

IHC: 25-100

Positive control: Human esophagus cancer

Predicted cell location: Cytoplasm

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Synthetic peptide of human LPP

**Formulation:** pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

**Concentration:** lot specific

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 66 kDa

Gene Name: LIM domain containing preferred translocation partner in lipoma

Database Link: NP 005569

Entrez Gene 4026 Human

Q93052

**Background:** This gene encodes a member of a subfamily of LIM domain proteins that are characterized by

an N-terminal proline-rich region and three C-terminal LIM domains. The encoded protein localizes to the cell periphery in focal adhesions and may be involved in cell-cell adhesion and

cell motility. This protein also shuttles through the nucleus and may function as a

transcriptional co-activator.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



**Synonyms:** DKFZp779O0231; FLJ30652; FLJ41512

## **Product images:**



Gel: 8%SDS-PAGE Lysate: 40 μg Lane: HepG2 cells


Primary antibody: TA351364 (LPP Antibody) at

dilution 1/350

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 20 seconds



Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using TA351364 (LPP Antibody) at dilution 1/30 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using TA351364 (LPP Antibody) at dilution 1/30, treated with synthetic peptide. (Original magnification: ×200)