

## **Product datasheet for TA351417**

## **MT-ND3 Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

Applications: IHC

Recommended Dilution: IHC: 25-100

Positive control: Human thyroid cancer Predicted cell location: Cytoplasm

Reactivity: Human
Host: Rabbit
Isotype: IgG

**Clonality:** Polyclonal

**Immunogen:** Synthetic peptide of human MT-ND3 **Formulation:** pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

**Concentration:** lot specific

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Background: NADH:ubiquinone oxidoreductase (complex I) is an extremely complicated multiprotein

complex located in the inner mitochondrial membrane. Human complex I is important for energy metabolism because its main function is to transport electrons from NADH to ubiquinone, which is accompanied by trans-location of protons from the mitochondrial matrix to the intermembrane space. Human complex I appears to consist of 41 subunits. A small number of complex I subunits are the products of mitochondrial genes (subunits 1-7),

while the remainder are nuclear encoded and imported from the cytoplasm. NADH

dehydrogenase subunit 3 (ND3) localizes to the hydrophobic protein fragment of complex I. Mutations in the gene encodiing for ND3 may be associated with Parkinson disease.




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Product images:**



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA351417 (MT-ND3 Antibody) at dilution 1/30 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA351417 (MT-ND3 Antibody) at dilution 1/30, treated with synthetic peptide. (Original magnification: ×200)