

Product datasheet for TA351483

CYP1A1 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB: 200-1000

WB positive control: Mouse lung tissue

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide of human CYP1A1

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 58 kDa

Gene Name: cytochrome P450 family 1 subfamily A member 1

Database Link: NP 000490

Entrez Gene 13076 MouseEntrez Gene 24296 RatEntrez Gene 1543 Human

P04798



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:


This gene, CYP1A1, encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and its expression is induced by some polycyclic aromatic hydrocarbons (PAHs), some of which are found in cigarette smoke. The enzyme's endogenous substrate is unknown; however, it is able to metabolize some PAHs to carcinogenic intermediates. The gene has been associated with lung cancer risk. A related family member, CYP1A2, is located approximately 25 kb away from CYP1A1 on chromosome 15.

Synonyms: AHH; AHRR; CP11; CYP1; CYPIA1; P1-450; P450-C; P450DX

Protein Families: Druggable Genome, P450, Transmembrane

Protein Pathways: Metabolism of xenobiotics by cytochrome P450, Retinol metabolism, Tryptophan metabolism

Product images:



Gel: 6%SDS-PAGE Lysate: 40 μg

Lane: Mouse lung tissue

Primary antibody: TA351483 (CYP1A1 Antibody)

at dilution 1/220

Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution

Exposure time: 30 seconds