

Product datasheet for TA351534S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Phospholipase C beta 3 (PLCB3) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 200-1000

WB positive control: PC3 and A431 cells, mouse lung tissue and Hela cells

IHC: 50-200

Positive control: Human prostate cancer

Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide of human PLCB3

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 139 kDa

Gene Name: phospholipase C beta 3

Database Link: NP 000923

Entrez Gene 18797 MouseEntrez Gene 29322 RatEntrez Gene 5331 Human

Q01970

Background: This gene encodes a member of the phosphoinositide phospholipase C beta enzyme family

that catalyze the production of the secondary messengers diacylglycerol and inositol 1,4,5-

triphosphate from phosphatidylinositol in G-protein-linked receptor-mediated signal

transduction. Alternative splicing results in multiple transcript variants.

Synonyms: FLJ37084

Protein Families: Druggable Genome







Protein Pathways:

Alzheimer's disease, Calcium signaling pathway, Chemokine signaling pathway, Gap junction, GnRH signaling pathway, Huntington's disease, Inositol phosphate metabolism, Long-term depression, Long-term potentiation, Melanogenesis, Metabolic pathways, Phosphatidylinositol signaling system, Vascular smooth muscle contraction, Wnt signaling pathway

Product images:




Gel: 6%SDS-PAGE Lysate: 40 µg Lane 1-4: PC3 cells A431 cells mouse lung tissue Hela cells Primary antibody: [TA351534] (PLCB3 Antibody) at dilution 1/200 Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution Exposure time: 3 minutes




Immunohistochemistry of paraffin-embedded Human prostate cancer tissue using [TA351534] (PLCB3 Antibody) at dilution 1/40 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human prostate cancer tissue using [TA351534] (PLCB3 Antibody) at dilution 1/40, treated with synthetic peptide. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human brain tissue using [TA351534] (PLCB3 Antibody) at dilution 1/40 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human brain tissue using [TA351534] (PLCB3 Antibody) at dilution 1/40, treated with synthetic peptide. (Original magnification: ×200)