

Product datasheet for TA351549S

POU6F2 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Human placenta tissue lysate

IHC: 50-100

Positive control: Human tonsil Predicted cell location: Nucleus

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide of human POU6F2

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 73 kDa

Gene Name: POU class 6 homeobox 2

Database Link: NP 009183

Entrez Gene 218030 MouseEntrez Gene 11281 Human

P78424

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US






Background:

This gene encodes a member of the POU protein family characterized by the presence of a bipartite DNA binding domain, consisting of a POU-specific domain and a homeodomain, separated by a variable polylinker. The DNA binding domain may bind to DNA as monomers or as homo- and/or heterodimers, in a sequence-specific manner. The POU family members are transcriptional regulators, many of which are known to control cell type-specific differentiation pathways. This gene is a tumor suppressor involved in Wilms tumor (WT) predisposition.

Synonyms: RPF-1; WT5; WTSL

Protein Families: Transcription Factors

Product images:



Gel: 8%SDS-PAGE Lysate: 40 μg

Lane: Human placenta tissue lysate


Primary antibody: [TA351549] (POU6F2 Antibody)

at dilution 1/800

Secondary antibody: Goat anti rabbit IgG at


1/5000 dilution

Exposure time: 3 minutes



Immunohistochemistry of paraffin-embedded Human tonsil tissue using [TA351549] (POU6F2 Antibody) at dilution 1/50 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human tonsil tissue using [TA351549] (POU6F2 Antibody) at dilution 1/50, treated with synthetic peptide. (Original magnification: ×200)