

Product datasheet for TA351639S

SDHB Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: PC3 and hepg2 cells, human fetal liver tissue

IHC: 25-100

Positive control: Human thyroid cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide of human SDHB

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 32 kDa

Gene Name: succinate dehydrogenase complex iron sulfur subunit B

Database Link: NP 002991

Entrez Gene 67680 MouseEntrez Gene 298596 RatEntrez Gene 6390 Human

P21912



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background: Complex II of the respiratory chain, which is specifically involved in the oxidation of succinate,

carries electrons from FADH to CoQ. The complex is composed of four nuclear-encoded subunits and is localized in the mitochondrial inner membrane. The iron-sulfur subunit is highly conserved and contains three cysteine-rich clusters which may comprise the iron-sulfur centers of the enzyme. Sporadic and familial mutations in this gene result in paragangliomas and pheochromocytoma, and support a link between mitochondrial

dysfunction and tumorigenesis.


Synonyms: CWS2; IP; PGL4; SDH; SDH1; SDH2; SDHIP

Protein Families: Druggable Genome

Protein Pathways: Alzheimer's disease, Citrate cycle (TCA cycle), Huntington's disease, Metabolic pathways,

Oxidative phosphorylation, Parkinson's disease

Product images:



Gel: 8%SDS-PAGE Lysate: 40 µg Lane 1-3: PC3 cells hepg2 cells

human fetal liver tissue


Primary antibody: [TA351639] (SDHB Antibody) at

dilution 1/400

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 30 seconds




Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA351639] (SDHB Antibody) at dilution 1/30 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA351639] (SDHB Antibody) at dilution 1/30, treated with synthetic peptide. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA351639] (SDHB Antibody) at dilution 1/30 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA351639] (SDHB Antibody) at dilution 1/30, treated with synthetic peptide. (Original magnification: ×200)