

Product datasheet for **TA351678S**


SLC16A10 Rabbit Polyclonal Antibody

Product data:


Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	WB: 500-2000 WB positive control: Mouse pancreas tissue lysate IHC: 25-50 Positive control: Human liver cancer Predicted cell location: Cell membrane
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Synthetic peptide of human SLC16A10
Formulation:	pH7.4 PBS, 0.05% NaN ₃ , 40% Glycerol
Purification:	Antigen affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	55 kDa
Gene Name:	solute carrier family 16 member 10
Database Link:	NP_061063 Entrez Gene 117247 Human Q8TF71
Background:	SLC16A10 is a member of a family of plasma membrane amino acid transporters that mediate the Na(+)-independent transport of aromatic amino acids across the plasma membrane. Sodium-independent transporter that mediates the uptake of aromatic acid. Can function as a net efflux pathway for aromatic amino acids in the basolateral epithelial cells.
Synonyms:	MCT10; PRO0813; TAT1
Protein Families:	Transmembrane

[View online »](#)


Product images:



Gel: 8%SDS-PAGE
 Lysate: 40 µg
 Lane: Mouse pancreas tissue lysate
 Primary antibody: [TA351678] (SLC16A10 Antibody) at dilution 1/600
 Secondary antibody: Goat anti rabbit IgG at 1/5000 dilution
 Exposure time: 3 minutes



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA351678] (SLC16A10 Antibody) at dilution 1/25 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA351678] (SLC16A10 Antibody) at dilution 1/25, treated with synthetic peptide. (Original magnification: ×200)