

## Product datasheet for **TA351784S**

### TAF10 Rabbit Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	WB: 500-2000 WB positive control: Human lung cancer tissue IHC: 25-100 Positive control: Human esophagus cancer Predicted cell location: Nucleus
Reactivity:	Human, Mouse
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Synthetic peptide of human TAF10
Formulation:	pH7.4 PBS, 0.05% NaN <sub>3</sub> , 40% Glycerol
Purification:	Antigen affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	22 kDa
Gene Name:	TATA-box binding protein associated factor 10
Database Link:	<a href="#">NP_006275</a> <a href="#">Entrez Gene 24075 Mouse</a> <a href="#">Entrez Gene 6881 Human</a> <a href="#">Q12962</a>



[View online »](#)

**Background:**

Initiation of transcription by RNA polymerase II requires the activities of more than 70 polypeptides. The protein that coordinates these activities is transcription factor IID (TFIID), which binds to the core promoter to position the polymerase properly, serves as the scaffold for assembly of the remainder of the transcription complex, and acts as a channel for regulatory signals. TFIID is composed of the TATA-binding protein (TBP) and a group of evolutionarily conserved proteins known as TBP-associated factors or TAFs. TAFs may participate in basal transcription, serve as coactivators, function in promoter recognition or modify general transcription factors (GTFs) to facilitate complex assembly and transcription initiation.

**Synonyms:**


TAF2A; TAF2H; TAFII30

**Protein Families:**

Druggable Genome, Transcription Factors

**Protein Pathways:**

Basal transcription factors

**Product images:**

Gel: 12%SDS-PAGE


Lysate: 40 µg

Lane: Human lung cancer tissue


Primary antibody: [TA351784] (TAF10 Antibody) at dilution 1/400

Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution


Exposure time: 10 seconds




Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using [TA351784] (TAF10 Antibody) at dilution 1/35 (Original magnification:  $\times 200$ )



Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using [TA351784] (TAF10 Antibody) at dilution 1/35, treated with synthetic peptide. (Original magnification:  $\times 200$ )



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA351784] (TAF10 Antibody) at dilution 1/35 (Original magnification:  $\times 200$ )



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA351784] (TAF10 Antibody) at dilution 1/35, treated with synthetic peptide. (Original magnification:  $\times 200$ )