

Product datasheet for TA351941S

PKM2 (PKM) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Jurkat, A549, hepg2 and NIH/3T3 cell

IHC: 25-100

Positive control: Human breast cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide of human PKM

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 58 kDa

Gene Name: pyruvate kinase, muscle

Database Link: NP 001193725

Entrez Gene 18746 MouseEntrez Gene 25630 RatEntrez Gene 5315 Human

P14618



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background: This gene encodes a protein involved in glycolysis. The encoded protein is a pyruvate kinase

that catalyzes the transfer of a phosphoryl group from phosphoenolpyruvate to ADP, generating ATP and pyruvate. This protein has been shown to interact with thyroid hormone and may mediate cellular metabolic effects induced by thyroid hormones. This protein has

been found to bind Opa protein, a bacterial outer membrane protein involved in gonococcal adherence to and invasion of human cells, suggesting a role of this protein in bacterial pathogenesis. Several alternatively spliced transcript variants encoding a few distinct

isoforms have been reported.


Synonyms: CTHBP; HEL-S-30; OIP3; PK3; PKM2; TCB; THBP1

Protein Families: Druggable Genome

Protein Pathways: Glycolysis / Gluconeogenesis, Metabolic pathways, Purine metabolism, Pyruvate metabolism,

Type II diabetes mellitus

Product images:



Gel: 8%SDS-PAGE Lysate: 40 μg Lane 1-4: Jurkat

A549

hepg2 and NIH/3T3 cell

Primary antibody: [TA351941] (PKM Antibody) at


dilution 1/300

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 10 seconds






Immunohistochemistry of paraffin-embedded Human breast cancer tissue using [TA351941] (PKM Antibody) at dilution 1/25 (Original magnification: ×200)




Immunohistochemistry of paraffin-embedded Human breast cancer tissue using [TA351941] (PKM Antibody) at dilution 1/25, treated with synthetic peptide. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA351941] (PKM Antibody) at dilution 1/25 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA351941] (PKM Antibody) at dilution 1/25, treated with synthetic peptide. (Original magnification: ×200)