

# **Product datasheet for TA351948**

### **WNT2 Rabbit Polyclonal Antibody**

#### **Product data:**

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Human endometrial cancer tissue

IHC: 30-150

Positive control: Human brain Predicted cell location: Cytoplasm

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

**Clonality:** Polyclonal

**Immunogen:** Synthetic peptide of human WNT2

**Formulation:** pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

**Concentration:** lot specific

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 40 kDa

**Gene Name:** Wnt family member 2

Database Link: NP 003382

Entrez Gene 22413 MouseEntrez Gene 7472 Human

P09544

**Background:** This gene is a member of the WNT gene family. The WNT gene family consists of structurally

related genes which encode secreted signaling proteins. These proteins have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and

patterning during embryogenesis. Alternatively spliced transcript variants have been

identified for this gene.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

### WNT2 Rabbit Polyclonal Antibody - TA351948

Synonyms: INT1L1; IRP


Protein Families: Adult stem cells, Cancer stem cells, ES Cell Differentiation/IPS, Secreted Protein, Stem cell

relevant signaling - Wnt Signaling pathway

**Protein Pathways:** Basal cell carcinoma, Hedgehog signaling pathway, Melanogenesis, Pathways in cancer, Wnt

signaling pathway

## **Product images:**



Gel: 8%SDS-PAGE Lysate: 40 μg


Lane: Human endometrial cancer tissue Primary antibody: TA351948 (WNT2 Antibody) at

dilution 1/300

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 10 seconds




Immunohistochemistry of paraffin-embedded Human brain tissue using TA351948 (WNT2 Antibody) at dilution 1/40 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human brain tissue using TA351948 (WNT2 Antibody) at dilution 1/40, treated with synthetic peptide. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human prostate cancer tissue using TA351948 (WNT2 Antibody) at dilution 1/40 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human prostate cancer tissue using TA351948 (WNT2 Antibody) at dilution 1/40, treated with synthetic peptide. (Original magnification: ×200)