

## **Product datasheet for TA352008S**

## **CLDND2 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Hela, 293T, PC3 and TM4 cell

IHC: 30-150

Positive control: Human breast cancer

Predicted cell location: Cytoplasm and Cell membrane

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Synthetic peptide of human CLDND2

**Formulation:** pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Gene Name:** claudin domain containing 2

Database Link: NP 689566

Entrez Gene 125875 Human

Q8NHS1



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

CLDND2 (claudin domain containing 2) is a 167 amino acid multi-pass membrane protein that belongs to the PMP-22/EMP/MP20 family and is encoded by a gene that maps to human chromosome 19q13.33. Consisting of around 63 million bases with over 1,400 genes, chromosome 19 makes up over 2% of human genomic DNA. Chromosome 19 includes a diversity of interesting genes and is recognized for having the greatest gene density of the human chromosomes. It is the genetic home for a number of immunoglobulin superfamily members including the killer cell and leukocyte lg-like receptors, a number of ICAMs, the CEACAM and PSG family, and Fc receptors. Key genes for eye color and hair color also map to chromosome 19. Peutz-Jeghers syndrome, spinocerebellar ataxia type 6, the stroke disorder CADASIL, hypercholesterolemia and insulin-dependent diabetes have been linked to chromosome 19.

Synonyms: MGC33839

**Protein Families:** Transmembrane

## **Product images:**



Gel: 12%SDS-PAGE Lysate: 40 µg Lane 1-4: Hela 293T PC3 and TM4 cell

Primary antibody: [TA352008] (CLDND2 Antibody)


at dilution 1/400

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 30 seconds





Immunohistochemistry of paraffin-embedded Human breast cancer tissue using [TA352008] (CLDND2 Antibody) at dilution 1/45 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human breast cancer tissue using [TA352008] (CLDND2 Antibody) at dilution 1/45, treated with synthetic peptide. (Original magnification: ×200)