

Product datasheet for **TA354654**

CTNND1 Rabbit Polyclonal Antibody

Product data:

| | |
|-------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Product Type: | Primary Antibodies |
| Applications: | Dot, WB |
| Recommended Dilution: | WB 0.1-1 µg/ml ELISA 0.01-0.1 µg/ml IP 2-5 µg/ml IHC 2-10 µg/ml FC 5-10 µg/ml |
| Reactivity: | Human, Mouse, Rat |
| Host: | Rabbit |
| Isotype: | IgG |
| Clonality: | Polyclonal |
| Immunogen: | A synthetic peptide corresponding to the epitope DNYGS at the phosphorylation site Tyr228 of human Catenin-delta1. |
| Formulation: | This affinity purified antibody is supplied in sterile Tris-buffered saline (pH7.2) containing antibody stabilizer. |
| Purification: | The Rabbit IgG is purified by site-modified Epitope Affinity Purification. |
| Conjugation: | Unconjugated |
| Storage: | Store at -20°C as received. |
| Stability: | Stable for 12 months from date of receipt. |
| Predicted Protein Size: | 88 kDa |
| Gene Name: | catenin delta 1 |
| Database Link: | NP_001078927 Entrez Gene 12388 Mouse Entrez Gene 311163 Rat Entrez Gene 1500 Human O60716 |



[View online »](#)


Background: Beta-catenin is a cytosolic, 88 kDa, 781 amino acid protein belongs to the β -catenin family. The N-terminus domain, containing the binding site and the phosphorylation sites. Beta-Catenin serves as a link between cytoskeleton actin and transmembrane cadherin(s). It is believed to contribute to tight cell-to-cell adhesion. It can enter the nucleus and interact with the TCF/LEF family of transcription factors, initiating gene expression. Normally, β -catenin transcriptional activity is suppressed by a Ser/Thr kinase termed GSK3 β and/or Casein Kinase I (CK1). Kinases are constitutively active and phosphorylates β -catenin at multiple sites, including S33 and S37, Y96, Y228, Y280 etc. Phosphorylation of β -catenin targets the molecule for degradation via a ubiquitination-mediated pathway. GSK3 β activity can be blocked by upstream signaling events such as Wnt-Frizzled interaction. This inhibits GSK3 β , allowing unphosphorylated β -catenin to enter the nucleus and initiate gene activation. The phosphorylation of beta-catenin might contribute to tumorigenesis.

Synonyms: CAS; CTNND; p120; p120(CAS); p120(CTN); P120CAS; P120CTN

Protein Families: Druggable Genome

Protein Pathways: Adherens junction, Leukocyte transendothelial migration

Product images:



WB: The cell lysate derived from HELA was immunoprobed at a dilution of 1:500 by the following antibodies: 1: Rabbit anti-beta-Catenin (pY228) 2: Rabbit anti-beta-Catenin (paired 228) 3: Rabbit anti-beta-Catenin (pY280) 4: Rabbit anti-beta-Catenin (Paired 280) 5: Negative control.



DB: 1 ug peptide was blot onto NC membrane A: Beta-Catenin (pY228) B: Beta-Catenin (Non-Phospho) C: Non-related Phosphopeptide Followed by immunoblotting Rabbit anti-beta-Catenin (pY228) at 1:1000.