

Product datasheet for **TA354685**

ERK2 (MAPK1) Rabbit Polyclonal Antibody

Product data:

| | |
|-------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Product Type: | Primary Antibodies |
| Applications: | WB |
| Recommended Dilution: | WB 0.1-1 µg/ml ELISA 0.01-0.1 µg/ml IP 2-5 µg/ml IHC 2-10 µg/ml FC 5-10 µg/ml |
| Reactivity: | Human, Mouse, Rat, Chicken, Canis |
| Host: | Rabbit |
| Isotype: | IgG |
| Clonality: | Polyclonal |
| Immunogen: | A synthetic peptide derived from epitope -LTEYV- without the dual phosphorylation sites Thr202 and Tyr204 of ERK1/2 protein from human, rat, mouse and dog origins |
| Formulation: | This affinity purified antibody is supplied in sterile Tris-buffered saline (pH7.2) containing antibody stabilizer. |
| Purification: | The Rabbit IgG is purified by Epitope Affinity Purification |
| Conjugation: | Unconjugated |
| Storage: | Store at -20°C as received. |
| Stability: | Stable for 12 months from date of receipt. |
| Predicted Protein Size: | 42/44 kDa |
| Gene Name: | mitogen-activated protein kinase 1 |
| Database Link: | NP_002736 Entrez Gene 26413 MouseEntrez Gene 116590 RatEntrez Gene 5594 Human P28482 |
| Background: | The extracellular signal-regulated kinases ½ (ERK1/2) are also known as MAPK (Mitogen-activated protein kinases) are a widely conserved family of serine/threonine protein kinases involved in many cellular programs such as cell proliferation, differentiation, motility, and death. The p44/42 MAPK (ERK1/2) signaling pathway can be activated in response to a diverse range of extracellular stimuli including mitogens, growth factors, and cytokines and is an important target in the diagnosis and treatment of cancer. The phosphorylation of MAPK at Thr202 and Tyr 204 are correlated to the progression and metastatic potential of tumor cells. |




[View online »](#)

Synonyms: ERK; ERK-2; ERK2; ERT1; MAPK2; p38; p40; p41; p41 mapk; p42-MAPK; P42MAPK; PRKM1; PRKM2

Protein Families: Druggable Genome, Protein Kinase

Protein Pathways: Acute myeloid leukemia, Adherens junction, Alzheimer's disease, Axon guidance, B cell receptor signaling pathway, Bladder cancer, Chemokine signaling pathway, Chronic myeloid leukemia, Colorectal cancer, Dorso-ventral axis formation, Endometrial cancer, ErbB signaling pathway, Fc epsilon RI signaling pathway, Fc gamma R-mediated phagocytosis, Focal adhesion, Gap junction, Glioma, GnRH signaling pathway, Insulin signaling pathway, Long-term depression, Long-term potentiation, MAPK signaling pathway, Melanogenesis, Melanoma, mTOR signaling pathway, Natural killer cell mediated cytotoxicity, Neurotrophin signaling pathway, NOD-like receptor signaling pathway, Non-small cell lung cancer, Oocyte meiosis, Pancreatic cancer, Pathways in cancer, Prion diseases, Progesterone-mediated oocyte maturation, Prostate cancer, Regulation of actin cytoskeleton, Renal cell carcinoma, T cell receptor signaling pathway, TGF-beta signaling pathway, Thyroid cancer, Toll-like receptor signaling pathway, Type II diabetes mellitus, Vascular smooth muscle contraction, VEGF signaling pathway

Product images:



WB: The whole cell lysate derived from 3T3 was separated in 10% SDS-PAGE, transferred onto NC membrane, and immunoblotted by Rabbit anti-ERK1/2 (Paired T202/Y204) antibody at 1:500. A dual immunoreactive band around ~42/44 kDa is observed.