

Product datasheet for TA354827

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

AMPK alpha 1 (PRKAA1) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB 0.1-1 μg/ml ELISA 0.01-0.1 μg/ml IP 2-5 μg/ml IHC 2-10 μg/ml FC 5-10 μg/ml

Reactivity: Human, Rat, Mouse, Bovine, Chicken, Dog, Invertebrate

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: A synthetic peptide surrounding to the epitope -LRTSC- with a phosphorylation site at Thr 172

of AMPK alpha protein from human origin

Formulation: This affinity purified antibody is supplied in sterile Tris-buffered saline (pH7.2) containing

antibody stabilizer.

Purification: The Rabbit IgG is purified by Epitope Affinity Purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 63 kDa

Gene Name: protein kinase AMP-activated catalytic subunit alpha 1

Database Link: NP 006242

Entrez Gene 65248 RatEntrez Gene 105787 MouseEntrez Gene 479351 DogEntrez Gene 5562

<u>Human</u> Q13131





Background:

AMP-activated protein kinase (AMPK) is highly conserved from yeast to plants and animals and plays a key role in the regulation of energy balance at both the cellular and the whole body levels. Once activated, it affects a metabolic switch from an anabolic to a catabolic state, both by acutely phosphorylating metabolic enzymes and, in the longer term, by regulating gene expression. AMPK is a heterotrimeric complex composed of a catalytic a subunit and regulatory β and ? subunits. Binding APM to the beta domains triggers increased phosphorylation at Thr172 on the activation loop of the alpha subunit. AMP Phosphorylation at Thr172 is catalyzed by the tumor suppressor kinase LKB1 or CaMKK-beta, TGF-beta activated kinase-1 (TAK1). AMPKa is also phosphorylated at Thr258 and Ser485 (for a1; Ser491 for a2).


Synonyms: AMPK; AMPKa1

Protein Families: Druggable Genome, Protein Kinase

Protein Pathways: Adipocytokine signaling pathway, Hypertrophic cardiomyopathy (HCM), Insulin signaling

pathway, mTOR signaling pathway, Regulation of autophagy

Product images:



WB: The whole cell lysate derived from HEK 293 was immunoblotted by Rabbit anti-AMPK alpha 1 antibody at 1:500. An immunereactive band is observed around ~63kDa.