

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA354828

JAK2 Rabbit Polyclonal Antibody

Product data:


Product Type:	Primary Antibodies
Applications:	Dot, WB
Recommended Dilution:	WB 0.1-1 μg/ml ELISA 0.01-0.1 μg/ml IP 2-5 μg/ml IHC 2-10 μg/ml FC 5-10 μg/ml
Reactivity:	Human
Host:	Rabbit
lsotype:	IgG
Clonality:	Polyclonal
Immunogen:	A synthetic peptide surrounding to the epitope -EYYK- with dual phosphorylation sites at Tyr1007 and Tyr1008 of JAK2 from human, mouse and rat origins
Formulation:	This affinity purified antibody is supplied in sterile Tris-buffered saline (pH7.2) containing antibody stabilizer.
Purification:	The Rabbit IgG is purified by site-modified Epitope Affinity Purification.
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	120 kDa (130 kDa)
Gene Name:	Janus kinase 2
Database Link:	<u>NP_001309127</u> <u>Entrez Gene 3717 Human</u> <u>O60674</u>
Background:	The Janus tyrosine kinase family is consists of four members, JAK1, JAK2, JAK3 and Tyk2. Upon the cytokine ligand binding to its the receptors, the dimerization of the receptors occurs followed by phosphorylation of the receptors. The phosphorylation of receptors recruites the tyrosine kinase physically associated to the dimerized receptors, and phosphorylate the kinase, thus activates the signal transducer and activator of thranscription (STAT1, 3, 5) and the other signal proteins.
Synonyms:	ЈТК10; ТНСҮТЗ
Protein Families:	Druggable Genome, Protein Kinase




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US Protein Pathways:

Adipocytokine signaling pathway, Chemokine signaling pathway, Jak-STAT signaling pathway

Product images:





WB: The tissue lysate derived from GH stimulated mouse liver was immunoblotted by Rabbit anti-JAK2 (pY1007/pY1008) antibody at 1:500.

DB: 1 ug peptide was blot onto NC membrane A: JAK2 (pY1008, pY1009) B: JAK2 (Non phosphospecific) C: Non-related Phosphopeptide) were blotted at a 1:2000 dilution by: 1: Rabbit anti-JAK2 (pY1008/pY1009), 2: Rabbit anti-JAK2 (Nonphospho specific).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US