

# **Product datasheet for TA354839**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# **Neurofibromin (NF1) Rabbit Polyclonal Antibody**

#### **Product data:**

**Product Type:** Primary Antibodies

**Applications:** Dot, WB

**Recommended Dilution:** WB 0.1-1 μg/ml ELISA 0.01-0.1 μg/ml IP 2-5 μg/ml IHC 2-10 μg/ml FC 5-10 μg/ml

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

**Immunogen:** A synthetic peptide surrounding the epitope –PTSPY- without phosphorylation at human

neurofibromin. This sequence is identical among human, rat, mouse, chicken, bovine and

dog.

Formulation: This affinity purified antibody is supplied in sterile Tris-buffered saline (pH7.2) containing

antibody stabilizer.

**Purification:** The Rabbit IgG is purified by Epitope Affinity Purification

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 240 kDa

**Gene Name:** neurofibromin 1

Database Link: NP 000258

Entrez Gene 4763 Human

P21359



#### Neurofibromin (NF1) Rabbit Polyclonal Antibody - TA354839

**Background:** The neurofibromatosis type 1 (NF1) tumor suppressor (neurofibromin) is thought to play

crucial roles in cellular Ras- and cAMP-dependent kinase (PKA)-associated signals. The absence of or alteration of the neurofibromin protein may lead to neurofibromatosis disease.

Recently, a cellular neurofibromin-associating protein, NG,NG-dimethylarginine

dimethylaminohydrolase (DDAH)has been identified as a cellular NO/NOS regulator which increases PKA phosphorylation of native neurofibromin in a dose-dependent manner. The PKA accessibility of neurofibromin regulated via DDAH interaction may modulate the cellular

function of neurofibromin that is implicated in NF1-related pathogenesis.

**Synonyms:** NFNS; VRNF; WSS


**Protein Families:** Druggable Genome

**Protein Pathways:** MAPK signaling pathway

# **Product images:**



WB: The whole cell lysates derived from hypoxic mouse brain were immunoblotted by Rabbit anti-NF1 (pSer2741) at 1:1000



DB: 1ug peptide was blot onto NC membrane A:NF1 (pS2741) B: NF1 (Nonphospho); C: Non-related phosphospecific PP were blotted at a 1:1000 dilution by: 1: Rabbit anti-NF1 (p2741); 2: Rabbit anti-NF1 (paired 2741)