

# **Product datasheet for TA354849**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Neuropilin 1 (NRP1) Rabbit Polyclonal Antibody

#### **Product data:**

**Product Type:** Primary Antibodies

**Applications:** Dot, WB

Recommended Dilution: WB 0.1-1 μg/ml ELISA 0.01-0.1 μg/ml IP 2-5 μg/ml IHC 2-10 μg/ml FC 5-10 μg/ml

Reactivity: Human, Bovine

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** A synthetic peptide surrounding the epitope –LNTQS- without phosphorylation site at Thr916.

This sequence is identical among human and bovine.

Formulation: This affinity purified antibody is supplied in sterile Tris-buffered saline (pH7.2) containing

antibody stabilizer.

**Purification:** The Rabbit IgG is purified by Epitope Affinity Purification

**Conjugation:** Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 130 kDa

Gene Name: neuropilin 1

Database Link: NP 001019799

Entrez Gene 8829 Human

<u>014786</u>

Background: The neuropilin-1 (Nrp1) is a multifunctional protein, identified principally as a receptor for the

class 3 semaphorins and members of the vascular endothelial growth factor (VEGF) family, but it is capable of other interactions. It is a marker of regulatory T cells (Tr), which often carry Nrp1 and latency-associated peptide (LAP)-TGF-beta1 (the latent form). Nrp1 is reported

to be expressed in breast cancer cells. The ischemia can increase neuropilin 1 protein

expression in experimental rat brain as well. The phosphorylation of Nrp1 is essential for the

Nrp1 activation.

Synonyms: BDCA4; CD304; NP1; NRP; VEGF165R






**Protein Families:** Druggable Genome, Secreted Protein, Transmembrane

**Protein Pathways:** Axon guidance

## **Product images:**



WB: The 4 mg protein of whole tissue lysates derived from human brain were immunoprecipitated by 4 ug of Rabbit antiNeuropilin1Paired T916), and probed by the same antibody at 1:1000. An immunoreactive band at ~130kDa was observed.



DB: 1 ug peptide was blot onto NC membrane A:Neuropilin1 (Phosphospecific Peptide at pT916) B: Neuropilin1 (Nonphospho PP); C: Non-related phosphospecific PP were blotted at a 1:1000 dilution by: 1: Rabbit anti-Neuropilin1 (pT916); 2: Rabbit anti-Neuropilin1 (paired T916)