

Product datasheet for **TA354861**

FUT4 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Recommended Dilution:	WB 0.1-1 µg/ml ELISA 0.01-0.1 µg/ml IP 2-5 µg/ml IHC 2-10 µg/ml FC 5-10 µg/ml
Reactivity:	Human
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	The Synthetic peptide corresponding to the C-term of human SSEA1 protein.
Formulation:	This affinity purified antibody is supplied in sterile Phosphate buffered saline (pH7.2) containing antibody stabilizer.
Purification:	The Rabbit IgG is purified by Epitope Affinity Purification
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	59 kDa
Gene Name:	fucosyltransferase 4
Database Link:	NP_002024 Entrez Gene 2526 Human P22083
Background:	CD15 is an antigen present on myelomonocytic cells and Lacto-Nfucopentose III (also known as hapten X). It is present on greater than 90% of granulocytes including neutrophils and eosinophils, and to a lesser degree, on monocytes. CD15 is expressed in Reed-Sternberg cells of Hodgkin's disease (of the nodular sclerosis, mixed cellularity and lymphocyte-depleted subtypes). This antibody recognizes myeloid cells, mainly granulocytes but not on B cells, T cells, monocytes, erythrocytes or platelets. Positive staining for CD15 combined with a negative reaction for lymphocytic markers may provide support for Hodgkin's disease. Fifty percent or more of adenocarcinomas have been described as showing significant cytoplasmic positivity for CD15. It does not detect most mesotheliomas.
Synonyms:	CD15; ELFT; FCT3A; FUC-TIV; FUTIV; LeX; SSEA-1



[View online »](#)

Protein Families: Transmembrane

Protein Pathways: Glycosphingolipid biosynthesis - lacto and neolacto series, Metabolic pathways