

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA354927

## LYRIC (MTDH) Rabbit Polyclonal Antibody

## **Product data:**

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB 0.1-1 μg/ml ELISA 0.01-0.1 μg/ml IP 2-5 μg/ml IHC 2-10 μg/ml FC 5-10 μg/ml
Reactivity:	Human, Mouse, Rat, Bovine
Host:	Rabbit
lsotype:	lgG
Clonality:	Polyclonal
Immunogen:	A synthetic peptide derived from the c-term of human AEG1. This sequence is identical to human, mouse, rat and bovine
Formulation:	This affinity purified antibody is supplied in sterile Tris-buffered saline (pH7.2) containing antibody stabilizer.
Purification:	The Rabbit IgG is purified by Epitope Affinity Purification
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	64 kDa
Gene Name:	metadherin
Database Link:	<u>NP_848927</u> <u>Entrez Gene 67154 MouseEntrez Gene 170910 RatEntrez Gene 92140 Human Q86UE4</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### **GRIGENE** LYRIC (MTDH) Rabbit Polyclonal Antibody – TA354927

Background:AEG1 (Astrocyte elevated gene 1), also called Metadherin (MTDH), or L LYsine-RIch CEACAM1<br/>co-isolated (LYRIC), is 582 as single transmembrane domain (TMD) protein, this protein is<br/>widely expressed and highly conserved between species. It is not a structural component for<br/>tight junctional (TJ) complexes, it is recruited during the maturation of the tight junction<br/>complex and co-localizes with tight junction proteins ZO-1 and occludin in polarized epithelial<br/>cells. This novel protein is also reported as an HIV-1 inducible gene, which is expressed<br/>ubiquitously with higher expression in tissues containing muscular actin and its expression is<br/>increased in astrocytes infected with HIV-1 or treated with gp120 or tumor necrosis factor<br/>(TNF)-alpha. AEG1 expression is elevated in subsets of breast carcinomas, malignant gliomas<br/>and melanomas. AEG1 may affect tumor progression in multiple cell lineages by augmenting<br/>expression of the transformed phenotype and/or by inducing glutamate excitotoxicity in<br/>malignant glioma. In these contexts, an HIV-1-inducible gene, AEG-1, may contribute to<br/>multiple brain abnormalities and tumor formation, by both common and distinct<br/>mechanisms.

Synonyms:	3D3; AEG-1; AEG1; LYRIC
Protein Families:	Transmembrane

### **Product images:**



WB: The cell lysate derived from Hela was immuno-blotted by Rabbit anti-AEG1 at 1:500. An immunoreactive band around 64kDa is observed.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US