

## Product datasheet for **TA355130**

### Lymphocyte Activation Gene 3 (LAG3) Mouse Monoclonal Antibody [Clone ID: 6D1]

#### Product data:


Product Type:	Primary Antibodies
Clone Name:	6D1
Applications:	ELISA, FC, IF, IHC
Recommended Dilution:	ICC start at 1 ug/mL. IF start at 10 ug/mL. IHC start at 5 ug/mL.
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1, kappa
Clonality:	Monoclonal
Immunogen:	LAG-3 antibody was raised against the extracellular domain of human LAG-3
Formulation:	LAG-3 Antibody is supplied in PBS containing 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	LAG-3 Antibody is supplied as protein A purified IgG1.
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	Predicted: 55 kDa; Observed: 60, 70, 80 kDa
Gene Name:	lymphocyte activating 3
Database Link:	<a href="#">NP_002277</a> <a href="#">Entrez Gene 3902 Human P18627</a>
Background:	The lymphocyte activation gene-3 (LAG3) is a member of the immunoglobulin superfamily and binds MHC class II with high affinity, negatively regulating T-cell function and homeostasis. It is expressed in B, T, and NK cells, monocytes, and dendritic cells, and acts to regulate T cell expansion. LAG3 is also an important immune checkpoint protein, with anti-LAG3 antibodies activating T effector cells and affecting regulatory T cell functions. Furthermore, LAG3 appears to act in a synergistic fashion with PD-1/PD-L1, suggesting that a dual antibody approach may prove useful in cancer immunotherapy.




[View online »](#)

Synonyms: CD223; FDC; LAG-3


Product images:




Titration curve analysis of LAG-3 mAbs to detect recombinant LAG-3 in ELISA with [TA355126], [TA355127], [TA355128], [TA355129], TA355130, [TA355131] and [TA355132] antibodies at decreasing concentrations.




Immunohistochemistry of LAG-3 in human lymphoma tissue using LAG-3 Antibody and control mouse IgG (corner box) at 5 ug/ml.




Immunocytochemistry of LAG-3 in over-expressing HEK293 cells using LAG-3 antibody and control mouse IgG antibody (left corner box) at 1 ug/ml.



Immunofluorescence of LAG-3 in over-expressing HEK293 cells using LAG-3 Antibody at 2 ug/ml.



Immunofluorescence of LAG-3 in over human spleen tissue using LAG-3 Antibody at 10 ug/ml.



Flow cytometry analysis of LAG-3 over expressing HEK293 cells using LAG-3 antibody at 1 ug/ml. Blue: untransfected HEK293 cells. Yellow: LAG-3 over expressing HEK293 cells.