

## Product datasheet for **TA355921**

### Pias2 Rabbit Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	WB
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	The immunogen is a synthetic peptide corresponding to a region of Mouse
Specificity:	<b>Expected reactivity:</b> Cow, Dog, Guinea Pig, Horse, Human, Mouse, Pig, Rabbit, Rat <b>Homology:</b> Cow: 86%; Dog: 86%; Guinea Pig: 86%; Horse: 93%; Human: 93%; Mouse: 100%; Pig: 93%; Rabbit: 79%; Rat: 93%
Formulation:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose. <i>Note that this product is shipped as lyophilized powder to China customers.</i>
Concentration:	lot specific
Purification:	Affinity Purified
Conjugation:	Unconjugated
Storage:	For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	68kDa
Gene Name:	protein inhibitor of activated STAT 2
Database Link:	<a href="#">NP_032628</a> <a href="#">Entrez Gene 17344 Mouse</a> <a href="#">Q8C5D8</a>




[View online »](#)

**Background:**

Pias2 functions as an E3-type small ubiquitin-like modifier (SUMO) ligase, stabilizing the interaction between UBE2I and the substrate, and as a SUMO-tethering factor. Pias2 plays a crucial role as a transcriptional coregulation in various cellular pathways, including the STAT pathway, the p53 pathway and the steroid hormone signaling pathway. The effects of this transcriptional coregulation, transactivation or silencing may vary depending upon the biological context and PIAS2 isoform studied. However, it seems to be mostly involved in gene silencing. Pias2 binds to sumoylated ELK1 and enhances its transcriptional activity by preventing recruitment of HDAC2 by ELK1, thus reversing SUMO-mediated repression of ELK1 transactivation activity. Isoform PIASx-beta, but not isoform PIASx-alpha, promotes MDM2 sumoylation. Isoform PIASx-alpha promotes PARK7 sumoylation. Isoform PIASx-beta promotes NCOA2 sumoylation more efficiently than isoform PIASx-alpha

**Synonyms:**

ARIP3; DIP; MGC102682; miz; MIZ1; PIASX; PIASX-ALPHA; PIASX-BETA; SIZ2; ZMIZ4

**Product images:**

WB Suggested Anti-Pias2 Antibody Titration: 0.2-1 ug/ml  
Positive Control: Mouse Thymus