

Product datasheet for TA356320

Fbxo2 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Reactivity: Mouse Rabbit Host:

Clonality: Polyclonal

Specificity: Expected reactivity: Cow, Dog, Guinea Pig, Horse, Human, Mouse, Rabbit, Rat, Zebrafish

Homology: Cow: 92%; Dog: 100%; Guinea Pig: 100%; Horse: 100%; Human: 100%; Mouse:

100%; Rabbit: 85%; Rat: 93%; Zebrafish: 77%

Formulation: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2%

sucrose.

Note that this product is shipped as lyophilized powder to China customers.

Concentration: lot specific

Purification: Affinity Purified Conjugation: Unconjugated

Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small

aliquots to prevent freeze-thaw cycles.

Shelf life: one year from despatch. Stability:

Predicted Protein Size: 34kDa

Gene Name: F-box protein 2 Database Link: NP 789818

Entrez Gene 230904 Mouse

Q80UW2



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

Fbxo2 is a substrate recognition component of a SCF (SKP1-CUL1-F-box protein) E3 ubiquitin-protein ligase complex that mediates the ubiquitination and subsequent proteasomal degradation of target proteins. Involved in the endoplasmic reticulum-associated degradation pathway (ERAD) for misfolded lumenal proteins by recognizing and binding sugar chains on unfolded glycoproteins that are retrotranslocated into the cytosol and promoting their ubiquitination and subsequent degradation. Prevents formation of cytosolic aggregates of unfolded glycoproteins that have been retrotranslocated into the cytosol. Able to recognize and bind denatured glycoproteins, preferentially those of the high-mannose type.

Synonyms: Fbg1; Fbs1; FBX2; NFB42

Product images:



WB Suggested Anti-Fbxo2 Antibody Titration: 1.0 ug/ml

Positive Control: Mouse Small Intestine