

## **Product datasheet for TA356502**

## **PCDH8 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type: Primary Antibodies** 

WB **Applications:** 

Reactivity: Human Rabbit Host:

Clonality: Polyclonal

Immunogen: The immunogen is a synthetic peptide directed towards the middle region of human PCDH8

Specificity: Expected reactivity: Cow, Dog, Guinea Pig, Horse, Human, Mouse, Rabbit, Rat

Homology: Cow: 100%; Dog: 100%; Guinea Pig: 100%; Horse: 93%; Human: 100%; Mouse:

100%; Rabbit: 100%; Rat: 100%

Formulation: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2%

sucrose.

Note that this product is shipped as lyophilized powder to China customers.

Concentration: lot specific

**Purification: Affinity Purified** Conjugation: Unconjugated

For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small Storage:

aliquots to prevent freeze-thaw cycles.

Shelf life: one year from despatch. Stability:

**Predicted Protein Size:** 101kDa

Gene Name: protocadherin 8

Database Link: NP 116567

Entrez Gene 5100 Human

Q5TAN1



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

This gene belongs to the protocadherin gene family, a subfamily of the cadherin superfamily. PCDH8 is an integral membrane protein that is thought to function in cell adhesion in a CNS-specific manner. Unlike classical cadherins, which are generally encoded by 15-17 exons, this gene includes only 3 exons. Notable is the large first exon encoding the extracellular region, including 6 cadherin domains and a transmembrane region. This gene belongs to the protocadherin gene family, a subfamily of the cadherin superfamily. The gene encodes an integral membrane protein that is thought to function in cell adhesion in a CNS-specific manner. Unlike classical cadherins, which are generally encoded by 15-17 exons, this gene includes only 3 exons. Notable is the large first exon encoding the extracellular region, including 6 cadherin domains and a transmembrane region. Alternative splicing yields isoforms with unique cytoplasmic tails.

Synonyms: ARCADLIN; PAPC
Protein Families: Transmembrane

## **Product images:**



WB Suggested Anti-PCDH8 Antibody Titration:

0.2-1 ug/ml

ELISA Titer: 1:12500

Positive Control: 721\_B cell lysate